

VALUATION EXTENSIONS OF FILTERED AND GRADED ALGEBRAS

C. BAETICA^{1,*} AND F. VAN OYSTAEYEN²

¹UNIVERSITY OF BUCHAREST, FACULTY OF MATHEMATICS, BUCHAREST,
ROMANIA

²UNIVERSITY OF ANTWERP, DEPARTMENT OF MATHEMATICS AND COMPUTER
SCIENCE, ANTWERP, BELGIUM

ABSTRACT. In this note we relate the valuations of the algebras appearing in the non-commutative geometry of quantized algebras to properties of sub-lattices in some vector spaces. We consider the case of algebras with *PBW*-bases and prove that under some mild assumptions the valuations of the ground field extend to a non-commutative valuation. Later we introduce the notion of *F*-reductor and graded reductor and reduce the problem of finding an extending non-commutative valuation to finding a reductor in an associated graded ring having a domain for its reduction.

Key Words: Filtered and graded reductors; Γ -separatedness; Valuation filtration.

2000 *Mathematics Subject Classification:* Primary 16A08.

INTRODUCTION

In the so-called non-commutative geometry of quantized algebras, some steps in the direction of a divisor theory via non-commutative valuation rings have already been taken; see Hussein and Van Oystaeyen (1996), Van Oystaeyen and Willaert (1996), Willaert (1997), etc. Most of the algebras appearing in the theory of rings of differential operators, quantized algebras of different kinds including many quantum groups, regular algebras in projective non-commutative geometry, come equipped with a natural gradation or filtration controlled by some finite dimensional vector space(s), e.g. the degree one part of filtration or gradation.

In this note we relate the valuations of the algebras considered to properties of sub-lattices in some vector space(s) as mentioned above. In the first section we generalize some results from Li and Van Oystaeyen (1990) to Γ -filtrations, where Γ is a totally ordered group. In Section 2 we consider the case of algebras with *PBW*-bases and prove that under some mild assumptions the valuations of the ground field extend to a non-commutative valuation. As an application we single out the case of Weyl algebras, although the results of this section apply to a large class of examples. In the last two sections we introduce the notion of *F*-reductor and

*Correspondence: C. Baetica, University of Bucharest, Faculty of Mathematics, Str. Academiei 14, RO-010014, Bucharest, Romania; E-mail: baetica@al.math.unibuc.ro.

graded reductor and reduce the problem of finding an extending non-commutative valuation to finding a reductor in an associated graded ring having a domain for its reduction.

Throughout this paper we use the notation in Van Oystaeyen (2000).

1. VALUATION FILTRATIONS

Let R be a ring and Γ a totally ordered group. A family FR of additive subgroups $F_\gamma R$, $\gamma \in \Gamma$ satisfying

- (i) $\gamma \leq \delta$ implies $F_\gamma \subseteq F_\delta$,
- (ii) $F_\gamma R F_\delta R \subseteq F_{\gamma+\delta} R$, for all $\gamma, \delta \in \Gamma$,
- (iii) $1 \in F_0 R$,

is a *filtration of type Γ* , or a Γ -filtration on R . For a Γ -filtration FR we may define the associated graded ring $G_F(R) = \bigoplus_{\gamma \in \Gamma} F_\gamma R / F_{<\gamma} R$, where $F_{<\gamma} R = \sum_{\gamma' < \gamma} F_{\gamma'} R$.

We say that FR is Γ -separated if for every $a \in R$, $a \neq 0$ there is a $\gamma \in \Gamma$ such that $a \in F_\gamma R - F_{<\gamma} R$. For $\Gamma = \mathbb{Z}$ the Γ -separatedness is equivalent to $\bigcap_{n \in \mathbb{Z}} F_n R = 0$, see Van Oystaeyen (2000, Remark 4.2.3), but for an arbitrary Γ the latter condition may be strictly weaker than this. If FR is Γ -separated, then we may define the *principal symbol map* $\sigma : R \rightarrow G_F(R)$ by $\sigma(a) = a \bmod F_{<\gamma} R$ whenever $a \in F_\gamma R - F_{<\gamma} R$. The *degree* $\deg \sigma(a) = \gamma$ of $\sigma(a)$ is uniquely determined. To FR one associates a value-function $v_F : R \rightarrow \Gamma \cup \{\infty\}$ defined by $v_F(0) = \infty$ and $v_F(a) = -\deg \sigma(a)$ for $a \neq 0$. It is well known that v_F is a valuation function on R whenever $G_F(R)$ is a domain; see Van Oystaeyen (2000, Corollary 4.2.7). Furthermore, we have that $1 \in F_0 R - F_{<0} R$, or equivalently $G_F(R)_0 \neq 0$.

All the filtrations we are dealing with are considered to be *exhaustive*, that is $\bigcup_{\gamma \in \Gamma} F_\gamma R = R$.

Definition 1.1. *If Δ is a skewfield and $\Lambda \subset \Delta$ is a subring, then Λ is a valuation ring of Δ if it is invariant under inner automorphisms of Δ and for $x \in \Delta$, $x \neq 0$, either $x \in \Lambda$ or $x^{-1} \in \Lambda$.*

Proposition 1.2. *Let R be an Artinian ring and FR a Γ -separated filtration on R . Then the following are equivalent:*

- (i) $G_F(R)$ is a domain,
- (ii) R is a skewfield and $G_F(R)$ is a gr-skewfield,
- (iii) R is a skewfield, $F_0 R$ is a valuation ring of R (with $F_{<0} R$ as unique maximal ideal), $\Gamma_s = \{\gamma \in \Gamma : G_F(R)_\gamma \neq 0\}$ is a subgroup of Γ and $G_F(R)$ is Γ_s -strongly graded.

Proof. (i) $\Rightarrow$ (ii) If $G_F(R)$ is a domain, then R is a domain; see Van Oystaeyen (2000, Proposition 4.2.4). Therefore R is a skewfield, because it is Artinian. For an element $\sigma(a_\gamma) \in G_F(R)_\gamma$, we have $a_\gamma \in F_\gamma R - F_{<\gamma} R$ and hence a_γ is invertible. Stand a_δ for its inverse. From the assumption $1 \in F_0 R - F_{<0} R$, it follows that $\gamma + \delta \geq 0$. Necessarily we get $\gamma + \delta = 0$, otherwise $\sigma(a_\gamma)\sigma(a_\delta) = 0$ contradicts the assumption that $G_F(R)$ is a domain. Thus $\delta = -\gamma$ and $\sigma(a_\delta)$ is an inverse of $\sigma(a_\gamma)$. (ii) $\Rightarrow$ (iii) Since $G_F(R)$ is a gr-skewfield, it is in particular a gr-domain. As Γ is a totally ordered group, $G_F(R)$ is a domain. By using again Van Oystaeyen (2000,

Proposition 4.2.4) we deduce that the value-function v_F associated to the filtration FR is a valuation function on R whose valuation ring is F_0R . The fact that Γ_s is a subgroup of Γ and $G_F(R)$ is Γ_s -strongly graded is obvious.

(iii) $\Rightarrow$ (i) $G_F(R)$ is strongly graded and $G_F(R)_0 = F_0R/F_{<0}R$ is a skewfield, so $G_F(R)$ is in particular a domain. $\square$

Remark 1.3. If FR is a Γ -separated *strong filtration*, i.e. $F_\gamma R F_\delta R = F_{\gamma+\delta} R$ for all $\gamma, \delta \in \Gamma$, then $\Gamma_s = \Gamma$. In particular, the valuation function v_F is surjective.

By Proposition 1.2 we get the following

Corollary 1.4. *If Δ is a skewfield and $\Lambda \subset \Delta$ is a subring, then the following are equivalent:*

- (i) Λ is a valuation ring of Δ ,
- (ii) There is a totally ordered group Γ and a valuation $v : \Delta \longrightarrow \Gamma \cup \{\infty\}$ such that Λ is the valuation ring of v ,
- (iii) There is a totally ordered group Γ and $F\Delta$ an exhaustive Γ -separated filtration on Δ such that $\Lambda = F_0\Delta$ and $G_F(\Delta)$ is a domain.

Proof. (i) $\Rightarrow$ (ii) See Van Geel (1981).

(ii) $\Rightarrow$ (iii) Set $F_\gamma\Delta = \{x \in \Delta : v(x) \geq -\gamma\}$, for $\gamma \in \Gamma$. Since v is surjective, the filtration $F\Delta$ is Γ -separated. Furthermore, $F\Delta$ is a strong filtration and thus $G_F(\Delta)$ is strongly graded with $G_F(\Delta)_0$ a skewfield. This is enough in order to show that $G_F(\Delta)$ is a domain.

(iii) $\Rightarrow$ (i) Easily enough. $\square$

Corollary 1.5. *Let FR be a Γ -separated filtration on a ring R . Then the following are equivalent:*

- (i) $G_F(R)$ is a gr-skewfield and $F_{<0}R \subseteq J(F_0R)$,
- (ii) R is a skewfield, F_0R is a valuation ring of R (with $F_{<0}R$ as unique maximal ideal), $\Gamma_s = \{\gamma \in \Gamma : G_F(R)_\gamma \neq 0\}$ is a subgroup of Γ and $G_F(R)$ is Γ_s -strongly graded.

Proof. (i) $\Rightarrow$ (ii) Let a be a non-zero element of R . Then there is $\gamma \in \Gamma$ such that $\sigma(a) \in G_F(R)_\gamma$. By hypothesis $G_F(R)$ is a gr-skewfield, hence there is $b \in R$ with $\sigma(a)\sigma(b) = 1$, that is $ab - 1 \in F_{<0}R$. It means that ab is invertible, hence a is also invertible. The other claims follow by Proposition 1.2.

(ii) $\Rightarrow$ (i) Readily by Proposition 1.2. $\square$

2. K -ALGEBRAS WITH PBW-BASES

From the preceding section we have seen how important is the Γ -separatedness for Γ -filtrations. A wide class of algebras are known to possess Γ -separated filtrations and among them we mention the K -algebras with a finite *PBW*-basis.

Throughout this section we consider K as being a field, Γ a totally ordered group and $v : K \longrightarrow \Gamma \cup \{\infty\}$ a valuation on K . We always assume that v is surjective, hence Γ is a commutative group. Set O_v the valuation ring of K associated to v , m_v its unique maximal ideal and $k_v = O_v/m_v$ the residue field. Then we define a Γ -filtration $f^v K$ on K by $f_\gamma^v K = \{x \in K : v(x) \geq -\gamma\}$ and we call it the *valuation filtration* on K . This is an exhaustive Γ -separated strong filtration.

Let A be an affine K -algebra generated by $a_1, \dots, a_n$, $K \langle \underline{X} \rangle$ the free K -algebra on the set $\underline{X} = \{X_1, \dots, X_n\}$ and $\pi : K \langle \underline{X} \rangle \rightarrow A$ the canonical K -algebras morphism given by $\pi(X_i) = a_i$, $i = 1, \dots, n$. Restriction of π to $O_v \langle \underline{X} \rangle$ defines a subring Λ of A , i.e. $\Lambda = \pi(O_v \langle \underline{X} \rangle)$. Denote by $\mathcal{R}$ the ideal of relations of A which is obviously a two-sided ideal. If $\mathcal{R}$ is generated by $p_1(\underline{X}), \dots, p_d(\underline{X})$ as a left ideal, then it is generated by $p_1(\underline{X}), \dots, p_d(\underline{X})$ as a two-sided ideal; see Hussein and Van Oystaeyen (1996, Lemma 2.1). Then we may assume that each $p_i(\underline{X}) \in O_v \langle \underline{X} \rangle$ (up to multiplying by a suitable constant), but not all are in $m_v \langle \underline{X} \rangle$. Since $\mathcal{R} \cap O_v \langle \underline{X} \rangle$ need not be generated again by the $p_1(\underline{X}), \dots, p_d(\underline{X})$ as a two-sided ideal of $O_v \langle \underline{X} \rangle$, we may encounter reduction problems. We say that Λ defines a *good reduction* of A (or that A *reduces well* at O_v) whenever $\overline{\mathcal{R}}$ is generated (as a two-sided ideal of $k_v \langle \underline{X} \rangle$) by the residues $\overline{p_i(\underline{X})}$, $i = 1, \dots, d$. For instance, if $\Lambda' = O_v \langle \underline{X} \rangle / \langle p_1(\underline{X}), \dots, p_d(\underline{X}) \rangle$ is a flat (equivalently, torsion-free) O_v -module, then Λ is a good reduction of A .

The subring Λ yields a filtration $F^\gamma A$ on A given by $F^\gamma A = (f^\gamma K)\Lambda$, $\gamma \in \Gamma$. We call $F^\gamma A$ the *valuation filtration* on A . Obviously, the valuation filtration $F^\gamma A$ is exhaustive. By $G_v(A)$ we denote the associated graded ring to the valuation filtration $F^\gamma A$.

The next result is Theorem 3.4.7 from Van Oystaeyen (2000).

Lemma 2.1. *For a graded K -algebra A that has a finite PBW-basis, the valuation filtration $F^\gamma A$ is Γ -separated and strong.*

This enables us to extend v to every graded K -algebra A that has a finite PBW-basis. In the following we denote by $A(k)$ the affine k -algebra defined by the same relations as A , where k is a suitable field.

Theorem 2.2. *Let A be a graded K -algebra with a finite PBW-basis and $\Lambda \subset A$ as before.*

- (i) *If $\overline{\Lambda} = \Lambda/m_v\Lambda$ is a domain and A is an Ore domain, then every valuation v on K extend to $Q = Q_d(A)$, the classical ring of fractions of A .*
- (ii) *If Λ defines a good reduction of A , $A(k_v)$ is a domain and A is an Ore domain, then every valuation v on K extend to Q .*

Proof. (i) The associated graded ring $G_v(A)$ is strongly graded with $G_v(A)_0 = \overline{\Lambda}$, hence $G_v(A)$ is a domain. By using Van Oystaeyen (2000, Corollary 4.2.7) we get a valuation function on A . It turns out that this is an extension of v to A and then we extend the valuation function on A to Q in the usual manner. All we have to do now is to show that $F^\gamma A \cap K = f^\gamma K$. It is clear that $f^\gamma K \subset F^\gamma A \cap K$. For converse, pick an element $x \in F^\gamma A \cap K$. Thus $x \in (f^\gamma K)\Lambda$ and $x \in K$, therefore $(f_{-\gamma}^\gamma K)x \subset \Lambda \cap K$. As $\Lambda \cap K = O_v$, we get $(f_{-\gamma}^\gamma K)(f_{-\gamma}^\gamma K)x \subset f_{-\gamma}^\gamma K$. But $(f_{-\gamma}^\gamma K)(f_{-\gamma}^\gamma K) = f_0^\gamma K = O_v$ and it is enough to conclude that $x \in f^\gamma K$.

(ii) Since Λ defines a good reduction of A , the ring $\overline{\Lambda}$ is isomorphic to $A(k_v)$, hence it is a domain. Now we can apply (i). $\square$

Corollary 2.3. *Let $D_n(K)$ be the skewfield of fractions of the Weyl algebra $A_n(K)$. Then every valuation v on K extends to $D_n(K)$.*

Proof. Set $A = A_n(K)$. Then A satisfies the conditions of Lemma 2.1 and thus the valuation filtration $F^v A$ is Γ -separated. The associated graded ring $G_v(A)$ is then strongly graded with $G_v(A)_0 = \Lambda/m_v \Lambda$, where Λ is defined as above. Note that Λ has also a finite *PBW*-basis over O_v . In particular Λ is O_v -flat, therefore it defines a good reduction of A . Consequently we can apply Proposition 2.2(ii) in order to get that every valuation v on K extend to $D_n(K)$. $\square$

For graded K -algebras with a finite *PBW*-basis the associated graded ring defined by the valuation filtration is a twisted group ring.

Proposition 2.4. *If A is a graded K -algebra that has a finite *PBW*-basis, then the associated graded ring $G_v(A)$ is isomorphic to the twisted group ring $\bar{\Lambda} * \Gamma$, where $\bar{\Lambda} = \Lambda/m_v \Lambda$.*

Proof. The Rees algebra $\tilde{A}$ of A associated to the valuation filtration $F^v A$ is defined by $\tilde{A} = \bigoplus_{\gamma \in \Gamma} F_\gamma^v A$. Since Λ is O_v -flat, we obtain

$$\tilde{A} = \bigoplus_{\gamma \in \Gamma} (f_\gamma K) \Lambda \simeq (\bigoplus_{\gamma \in \Gamma} f_\gamma K) \otimes_{O_v} \Lambda = \tilde{K} \otimes_{O_v} \Lambda.$$

From $G_v(A) \simeq \tilde{A}/\tilde{A}\Gamma^+$ we get that $G_v(A) \simeq \tilde{K}/\tilde{K}\Gamma^+ \otimes_{O_v} \Lambda \simeq G_v(K) \otimes_{O_v} \Lambda$. On the other side, $G_v(K)$ is a commutative strongly graded ring with the trivial action, therefore it is isomorphic to the twisted group ring $k_v * \Gamma$ and thus we get $G_v(A) \simeq (k_v * \Gamma) \otimes_{O_v} \Lambda \simeq \Lambda/m_v \Lambda * \Gamma$. $\square$

In particular, for Weyl algebras we obtain

Corollary 2.5. $G_v(A_n(K)) \simeq A_n(k_v) * \Gamma$.

For a plenty of examples of algebras with a finite *PBW*-basis, the reader is referred to Berger (1992) and Van Oystaeyen (2000, Observation 4.3.8).

Similarly to Hussein and Van Oystaeyen (1996, Proposition 2.3) we can prove the following

Proposition 2.6. *Let A be a graded K -algebra with a finite *PBW*-basis and $\Lambda \subset A$ as before. If A is a prime Goldie ring and $\bar{\Lambda}$ is a domain, then*

- (i) $Q = Q_{cl}(A)$ is a skewfield and $F^v A$ extends to a strong filtration on Q such that $F_0^v Q$ is a valuation ring of Q containing O_v , that is v extends to Q .
- (ii) $Q' = Q^g(A)$, the graded ring of fractions of A , is a gr-skewfield and $F^v Q$ induces a graded valuation on Q' .

3. FILTERED AND GRADED REDUCTORS

Let A be a K -algebra, K a field and FA an exhaustive separated $\mathbb{Z}$ -filtration such that $K \subseteq F_0 A$. Suppose in addition that FA is *finite*, i.e. $\dim_K F_n A < \infty$ for all $n \in \mathbb{Z}$. Since FA is finite and separated, it must be *left limited*, i.e. there is $n_0 \in \mathbb{Z}$ such that $F_n A = 0$ for all $n \leq n_0$. Without loss of generality we may suppose that the filtration FA is *positive*, that is $F_n A = 0$ for all $n < 0$.

Let v be a valuation on K with the value group Γ , $O_v \subset K$ the corresponding valuation ring, m_v its unique maximal ideal and $k_v = O_v/m_v$ the residue field.

Definition 3.1. Let V be a finite dimensional K -vector space and M an O_v -submodule of V . Then M is an O_v -lattice in V if it contains a K -basis of V and is a submodule of a finitely generated O_v -submodule of V . Usually, we will denote the k_v -vector space $M/m_v M$ by $\overline{V}$.

Remark 3.2. For any O_v -lattice M in V we have $\dim_{k_v} \overline{V} \leq \dim_K V$. When equality holds we say that M defines an *unramified reduction* of V .

The next result deals with some elementary properties of O_v -lattices that we did not find explicitly in the literature.

Lemma 3.3. Let V be a finite dimensional K -vector space, $V' \subset V$ a K -vector subspace and $M \subset V$ an O_v -submodule.

- (i) If M is an O_v -lattice in V , then the quotient module $M/M \cap V'$ is an O_v -lattice in V/V' .
- (ii) If $M/M \cap V'$ is an O_v -lattice in V/V' and $M \cap V'$ is an O_v -lattice in V' , then M is an O_v -lattice in V .

Proof. (i) It is known that $M \cap V'$ is an O_v -lattice in V' ; see Fossum (1973, Proposition 2.2(ii)). Then it is straightforward that $M/M \cap V'$ is an O_v -lattice in V/V' .

(ii) Easy. $\square$

The *valuation filtration* $F^\gamma V$ on V defined by $F^\gamma V = (f_\gamma^v K)M$ is an exhaustive filtration, but we do not know if it is Γ -separated or not. It is rather easy to see that the valuation filtration is Γ -separated whenever M is a free O_v -module. This happens, for instance, when $\Gamma = \mathbb{Z}$. This shows that the discrete case can be handled with ease.

Definition 3.4. Let A be a filtered K -algebra with a finite filtration FA and $\Lambda \subset A$ a subring. Then Λ is an F -reductor if $\Lambda \cap K = O_v$ and $\Lambda \cap F_n A$ is an O_v -lattice in $F_n A$ for all $n \in \mathbb{N}$.

We call the ring $\overline{A} = \Lambda/m_v \Lambda$ the (filtered) *reduction* of A with respect to Λ . The *valuation filtration* $F^\gamma A$ on A is defined by $F^\gamma A = (f_\gamma^v K)\Lambda$, $\gamma \in \Gamma$, while the *induced filtration* $F\Lambda$ of FA on Λ is given by $F_n \Lambda = \Lambda \cap F_n A$, $n \in \mathbb{N}$.

Proposition 3.5. Let A be as before and $\Lambda \subset A$ an F -reductor. We have

- (i) $A = K\Lambda$ and the valuation filtration $F^\gamma A$ is exhaustive,
- (ii) $F^\gamma A$ is Γ -separated, provided that $F^\gamma(F_n A)$ is Γ -separated for all $n \in \mathbb{N}$,
- (iii) the induced filtration of $F^\gamma A$ on $F_n A$ is the valuation filtration on $F_n A$ and provided that it is Γ -separated we get $G_v(F_n A) = \overline{F_n A} * \Gamma$, where $\overline{F_n A} = F_n \Lambda / m_v F_n \Lambda$,
- (iv) $(\overline{F_n A})_{n \in \mathbb{N}}$ defines a finite filtration on $\overline{A}$, denoted by $\overline{F} \overline{A}$ and $G_{\overline{F}}(\overline{A}) = \overline{G(A)}$, where $\overline{G(A)} = G(\Lambda)/m_v G(\Lambda)$.

Proof. (i) Since $F_n \Lambda$ is an O_v -lattice in $F_n A$, we have $K(F_n \Lambda) = F_n A$ for all $n \in \mathbb{N}$. As FA is exhaustive we get $K\Lambda = A$. The valuation filtration $f^v K$ is exhaustive and $K\Lambda = A$, therefore $F^\gamma A$ is an exhaustive filtration.

(ii) In order to show that $F^\gamma A$ is Γ -separated we first note that

$$(f_\gamma^v K)\Lambda \cap F_n A = (f_\gamma^v K)(\Lambda \cap F_n A) \quad (1)$$

for all $\gamma \in \Gamma$ and $n \in \mathbb{N}$. It is obvious that $(f_\gamma^v K)(\Lambda \cap F_n A) \subseteq (f_\gamma^v K)\Lambda \cap F_n A$. The converse inclusion follows readily if one observes that the filtration $f^v K$ is strong. To enter the detail, pick an element $x \in (f_\gamma^v K)\Lambda \cap F_n A$. It follows that $(f_{-\gamma}^v K)x \subseteq \Lambda \cap F_n A$ and therefore $x \in (f_\gamma^v K)(\Lambda \cap F_n A)$.

For any element $x \in A$, $x \neq 0$, there exists an $n \in \mathbb{N}$ such that $x \in F_n A$. Since the valuation filtration on $F_n A$ is assumed to be Γ -separated, there exists an element $\gamma \in \Gamma$ such that $x \in F_\gamma^v(F_n A) - F_{<\gamma}^v(F_n A)$. By using (1) we obtain that $x \in F_\gamma^v A - F_{<\gamma}^v A$.

(iii) By using (1) again we get that the induced filtration of $F^v A$ on $F_n A$ is the valuation filtration on $F_n A$. The crossed product structure of $G_v(F_n A)$ can be seen in the same way as in the proof of Proposition 2.4.

(iv) Let us remark that $m_v(\Lambda \cap F_n A) = m_v \Lambda \cap F_n A$. This can also be obtained by using (1). Then $(\overline{F_n A})_{n \in \mathbb{N}}$ defines a finite filtration on $\overline{A}$; see Remark 3.2. It is easily seen that $G_{\overline{F}}(\overline{A}) = \overline{G(A)}$. $\square$

It is a common strategy to deduce properties of A from properties of $G_F(A)$ whenever possible, so let us focus on the graded situation for a moment.

Definition 3.6. Let R be an $\mathbb{N}$ -graded K -algebra with $K \subseteq R_0$. Suppose that R is locally finite, i.e. $\dim_K R_n < \infty$ for all $n \in \mathbb{N}$ and let $\Lambda \subset R$ be a graded subring. Then Λ is called a graded reductor if $\Lambda \cap K = O_v$ and $\Lambda \cap R_n$ is an O_v -lattice in R_n for all $n \in \mathbb{N}$.

The ring $\overline{R} = \Lambda/m_v \Lambda$ is called the (graded) reduction of R with respect to Λ . The valuation filtration $F^v R$ on R is similarly defined by $F_\gamma^v R = (f_\gamma^v K)\Lambda$, $\gamma \in \Gamma$. A Γ -filtration FR on R such that $F_\gamma R = \bigoplus_{n \in \mathbb{N}} F_\gamma R \cap R_p$, for all $\gamma \in \Gamma$ is called a graded filtration.

Proposition 3.7. Let R be as before and $\Lambda \subset R$ a graded reductor. We have

- (i) $R = K\Lambda$ and the valuation filtration $F^v R$ is an exhaustive graded filtration,
- (ii) $F^v R$ is Γ -separated, provided that $F^v R_n$ is Γ -separated for all $n \in \mathbb{N}$,
- (iii) the induced filtration of $F^v R$ on R_n is the valuation filtration on R_n and provided that it is Γ -separated we get $G_v(R_n) = \overline{R}_n * \Gamma$, where $\overline{R}_n = \Lambda_n/m_v \Lambda_n$,
- (iv) $\overline{R} = \bigoplus_n \overline{R}_n$ and $G_v(R) = \overline{R} * \Gamma$.

Proof. For (i) and (iii) we can argue similarly to the proof of Proposition 3.5, while (iv) is rather obvious.

(ii) Pick an element $x \in R$, $x \neq 0$ and write $x = \sum_{i=0}^n x_i$ with $x_i \in R_i$ not all zero. For each x_i with $x_i \neq 0$ we get an element $\gamma_i \in \Gamma$ such that $x_i \in F_{\gamma_i}^v R_i - F_{<\gamma_i}^v R_i$. Assume that γ is the maximum of the γ_i . We claim that $x \in F_\gamma^v R - F_{<\gamma}^v R$. If $x \in F_{<\gamma}^v R$, then there is $\delta \in \Gamma$, $\delta < \gamma$, with the property that $x \in F_\delta^v R$. Since $F_\delta^v R = (f_\delta^v K)\Lambda$, we can write $x = \sum_{j=1}^s \alpha_j \lambda_j$, where $\alpha_j \in f_\delta^v K$ and $\lambda_j \in \Lambda$. Each λ_j can be written as follows $\lambda_j = \sum_n \lambda_{jn}$, with $\lambda_{jn} \in \Lambda_n$, and thus we get $x = \sum_n (\sum_{j=1}^s \alpha_j \lambda_{jn})$. In particular, $x_i = \sum_{j=1}^s \alpha_j \lambda_{ji}$ and therefore $x_i \in (f_\delta^v K)\Lambda_i$, a contradiction. $\square$

Note that the reductor property for Λ is "almost" sufficient to make $F^v R$ into a Γ -separated filtration, whereas in Hussein and Van Oystaeyen (1996) this was

obtained (for $\Gamma = \mathbb{Z}$) by restricting to *connected* positively graded K -algebras, that is, $R = K \oplus R_1 \oplus \cdots \oplus R_n \oplus \cdots = K[R_1]$ and $\dim_K R_1 < \infty$.

Lemma 3.8. *If Λ is a graded reductor for R and S is an Ore set in R , then there is an Ore set S' in Λ such that $(S')^{-1}\Lambda = S^{-1}R$.*

Proof. Take $S' = K^\times S \cap \Lambda$, where $K^\times = K - \{0\}$. Since $K\Lambda = R$, for every finite subset $\{a_1, \dots, a_m\}$ of R there exists a nonzero $\lambda \in O_v$ such that $\lambda a_i \in \Lambda$ for all $i = 1, \dots, m$. It is straightforward to check the Ore condition for S' and moreover $(S')^{-1}\Lambda = S^{-1}R$ is clear as well. $\square$

Corollary 3.9. *Suppose that $\Gamma = \mathbb{Z}$ and $G_v(R)$ is a domain. Then the localized filtration on $S^{-1}R$ derived from $F^v R$, denoted by $F^v S^{-1}R$, is exactly the localized filtration deriving from $m_v \Lambda$ -adic filtration on Λ .*

Similarly to Hussein and Van Oystaeyen (1996, Proposition 2.3) we have the following

Proposition 3.10. *Let R be as before and $\Lambda \subset R$ a graded reductor such that $F^v R$ is Γ -separated. Suppose that R is a prime Goldie ring and $\overline{R}$ is a domain. Then*

- (i) $Q = Q_{cl}(R)$ is a skewfield and $F^v R$ extends to a strong filtration on Q such that $F_0^v Q$ is a valuation ring of Q containing O_v , that is v extends to Q .
- (ii) $Q' = Q^g(R)$, the graded ring of fractions of R , is a gr-skewfield and $F^v Q$ induces a graded valuation on Q' .

For connected positively graded K -algebras, as are all the graded algebras appearing in the non-commutative geometry of *Proj*, the existence of graded reducers can be expressed in terms of certain unramifiedness properties.

Let R be a connected positively graded K -algebra with $\dim R_1 = n$, $\underline{X} = \{X_1, \dots, X_n\}$, and take $\pi : K \langle \underline{X} \rangle \rightarrow R$ a presentation of R . If we set $\Lambda = \pi(O_v \langle \underline{X} \rangle)$, then $\dim_{k_v} \overline{R}_1 = n$ since no elements of degree one in the gradation of $K \langle \underline{X} \rangle$ are in $\mathcal{R}$, the ideal of relations of R . Nevertheless, $\dim_K R_n$ and $\dim_{k_v} \overline{R}_n$ may be different for $n > 1$.

Proposition 3.11. *Let R be a connected positively graded K -algebra.*

- (i) *If $\dim_K R_n = \dim_{k_v} \overline{R}_n$ for all $n \in \mathbb{N}$, then Λ is a graded reductor of R .*
- (ii) *For $\Gamma = \mathbb{Z}$ the converse of (i) holds.*
- (iii) *When moreover $\overline{R}$ is a Goldie domain, we have that R is a domain and the $m_v \Lambda$ -adic filtration on R is induced by a valuation filtration on the skewfield of microfractions of R .*

Proof. (i) We have to prove that $\Lambda_n \subset R_n$ is an O_v -lattice for all $n \in \mathbb{N}$. Take $\{\overline{x}_1, \dots, \overline{x}_m\}$ a k_v -basis of $\overline{R}_n$. Then the elements $x_1, \dots, x_m$ are linearly independent over K and by hypothesis they form a K -basis of R_n . This shows that Λ_n contains a K -basis of R_n . As Λ_n is a finitely generated O_v -module we can conclude that Λ_n is an O_v -lattice in R_n .

(ii) When $\Gamma = \mathbb{Z}$, each graded component Λ_n of Λ is a free O_v -module whose rank equals $\dim_K R_n$, as long as Λ is a graded reductor of R .

(iii) In this case $m_v = \pi O_v$, $\pi \in O_v$ and $F_n^v R = \pi^{-n} \Lambda$ for all $n \in \mathbb{Z}$. $F^v R$ is a strong filtration, hence $G_v(R)$ is strongly graded and its degree zero component $G_v(R)_0$

equals $\overline{R}$ which is a domain. Thus $G_v(R)$ is a domain, hence R is a domain too. Furthermore, in this case we have $G_v(R) = \overline{R}[t, t^{-1}]$. Since $\overline{R}$ is a Goldie domain, $\overline{R}[t, t^{-1}]$ is also a Goldie domain and it has a skewfield of fractions $\overline{\Delta}$ as well as a graded skewfield of fractions $\overline{\Delta}^g$. The multiplicative set $R - \{0\}$ has $\sigma(R - \{0\}) = G_v(R) - \{0\}$ which is an Ore set because $G_v(R)$ is a Goldie domain. Setting $S_0 = \Lambda - \{0\}$ we get $\sigma(S_0) = G_v(R)_{\leq 0} - \{0\}$, where $G_v(R)_{\leq 0} = \bigoplus_{n \leq 0} G_v(R)_n = G_v(\Lambda)$. Clearly $\sigma(S_0)$ is an Ore set in $G_v(\Lambda)$ and $\sigma(S_0)^{-1}G_v(\Lambda) = Q^g(G_v(R))$.

It follows easily that for every $p \geq 1$, S maps to an Ore set $S^{(p)}$ of $\Lambda/m_v^p\Lambda$. Given $s \in S_0$, $a \in R$, the left Ore condition for $\sigma(S_0)$ yields an $s' \in S_0$, $a' \in R$, such that $s'a - a's \in m_v\Lambda$, say $s'a - a's = \pi^m b$ for some $b \in \Lambda$, $m \in \mathbb{Z}$ (here π is a generator of m_v). Now $s''b - a''s \in m_v\Lambda$ yields $(s''s')a - (s''a')s = \pi^m(a''s - \pi^{m'}y)$ for some $y \in \Lambda$, $m' \neq 0$. Hence $(s''s')a - (s''a' + \pi^m a'')s = \pi^{m+m'}y$. Consequently the microlocalization $Q^\mu(\Lambda)$ of Λ at S_0 may be defined by $Q^\mu(\Lambda) = \varprojlim S^{(p)}(\Lambda/m_v^p\Lambda)$. Obviously $Q^\mu(R) = Q^\mu(\Lambda)$ is a skewfield and it has a filtration whose associated graded ring is $\overline{\Delta}^g$, which is a domain and a graded skewfield. Now we can apply Proposition 3.10 and conclude that the filtration on $Q^\mu(R)$ is a valuation filtration. $\square$

4. LIFTING FROM GRADED TO FILTERED RINGS

Let A be a filtered K -algebra with a finite filtration FA and $\Lambda \subset A$ a subring. Then FA induces a filtration $F\Lambda$ on Λ such that $G_F(\Lambda) \subset G_F(A)$. The next result shows that there is a strong connection between graded and F -reductors.

Proposition 4.1. *Let A be as before and $\Lambda \subset A$ a subring.*

- (i) *If $\Lambda \subset A$ is an F -reductor, then $G_F(\Lambda) \subset G_F(A)$ and $\tilde{\Lambda} \subset \tilde{A}$ are graded reductors.*
- (ii) *If $G_F(\Lambda) \subset G_F(A)$ or $\tilde{\Lambda} \subset \tilde{A}$ are graded reductors, then $\Lambda \subset A$ is an F -reductor.*

Proof. (i) In order to show that $G_F(\Lambda) \subset G_F(A)$ is a graded reductor we use Lemma 3.3(i). That $\tilde{\Lambda} \subset \tilde{A}$ is a graded reductor follows immediately by definition.

(ii) By induction using Lemma 3.3(ii). $\square$

Consequently any F -reductor give rise to a graded reductor. On the other side, a graded reductor is an F -reductor, where FR is the grading filtration.

While the finite dimensional K -algebras have always a reductor, see Van Oystaeyen (1975, Proposition 54 and Theorem 56), in the infinite dimensional case the construction of reductors is not always easy.

Examples 4.2. (i) Consider g a finite dimensional Lie algebra over K and $A = U(g)$ the enveloping algebra of g . We may define a finite dimensional Lie algebra g_{O_v} over O_v with the same basis and the induced bracket. Let us fix a K -basis $\{x_1, \dots, x_n\}$ for g . We have structure constants $\lambda_{ij}^k \in K$ with $[x_i, x_j] = \sum_{k=1}^n \lambda_{ij}^k x_k$. Without loss of generality we may assume that $\lambda_{ij}^k \in O_v$ (up to multiplying all x_i by a suitable constant in O_v) but not all in m_v . Set $g_{O_v} = O_v x_1 + \dots + O_v x_n$. This is a Lie O_v -algebra with the induced bracket. Furthermore, g_{O_v} is an O_v -lattice

in g . On $\bar{g} = g_{O_v}/m_v g_{O_v}$ we define a Lie algebra structure over k_v by setting $[\bar{x}_i, \bar{x}_j] = \sum_{k=1}^n \bar{\lambda}_{ij}^k \bar{x}_k$, where the $\bar{x}_i$ are the images of the x_i in g_{O_v} and $\bar{\lambda}_{ij}^k$ are the images of λ_{ij}^k in k_v . By our assumptions $\bar{g}$ is not the trivial Lie algebra. Of course, $\bar{g}$ depends on the choice of the K -basis in g .

Let $\Lambda = U_{O_v}(g_{O_v})$ be the enveloping algebra of g_{O_v} . Consider on A the standard filtration FA and on Λ the induced filtration $F\Lambda$. We have that the filtration FA is finite, $G_F(\Lambda) = O_v[X_1, \dots, X_n]$ is a subring of the polynomial ring $G_F(A) = K[X_1, \dots, X_n]$ and obviously it is a graded reductor. Proposition 4.1(ii) shows that Λ is an F -reductor. Furthermore, the filtration $F^v A$ is Γ -separated, $G_v(A)$ is a domain isomorphic to $U_{k_v}(\bar{g})$ and thus we can extend v to $D(g) = Q_{cl}(U(g))$.

(ii) For the Weyl algebra $R = A_n(K)$ we take $\Lambda = A_n(O_v)$. We claim that Λ is a graded reductor of R . First note that Λ is a free O_v -module, therefore it defines a good reduction of R . On the other side, we have that $\text{rank}_{O_v} \Lambda_n = \dim_K R_n$ for all $n \in \mathbb{N}$, since Λ and R have the same PBW-basis, and so Λ is a graded reductor of R . It is also an F -reductor since the associated graded ring of $A_n(K)$ with respect to the Bernstein filtration is a polynomial ring.

On $G_v(A)$ we define a filtration $fG_v(A)$ by putting $f_n G_v(A) = G_v(F_n A)$, $n \in \mathbb{N}$. When $G_F(\Lambda) \subset G_F(A)$ is a graded reductor we have a valuation filtration $f^v G_F(A)$ on $G_F(A)$ given by $f_\gamma^v G_F(A) = (f_\gamma^v K)G_F(\Lambda)$, $\gamma \in \Gamma$. As in the proof of Proposition 3.7 we can show that the valuation filtration $f^v G_F(A)$ on $G_F(A)$ is Γ -separated, provided that $F^v A$ is Γ -separated.

Corollary 4.3. *Let A be as before and $\Lambda \subset A$ an F -reductor such that the filtration $F^v A$ is Γ -separated. Then $G_v(G_F(A))$ is isomorphic to $\overline{G(A)} * \Gamma$ and $G_f(G_v(A)) = G_v(G_F(A))$.*

Proof. We have $G_v(G_F(A)) = \bigoplus_{\gamma \in \Gamma} (f_\gamma^v K)G_F(\Lambda) / (f_{<\gamma}^v K)G_F(\Lambda)$ and since $G_F(\Lambda)$ is O_v -flat we get $G_v(G_F(A)) = \bigoplus_{\gamma \in \Gamma} (f_\gamma^v K) / (f_{<\gamma}^v K) \otimes_{O_v} G_F(\Lambda)$. The latter is nothing else but $G_v(K) \otimes_{O_v} G_F(\Lambda) = (k_v * \Gamma) \otimes_{O_v} G_F(\Lambda) = \overline{G(A)} * \Gamma$.

The fact that $G_f(G_v(A))$ equals $G_v(G_F(A))$ follows by the general compatibility result of Hussein and Van Oystaeyen (1996, Proposition 2.4) or can be verified directly. $\square$

Proposition 4.4. *Let A be as before and $\Lambda \subset A$ an F -reductor such that $F^v R$ is Γ -separated. If $\overline{G(A)}$ is a domain, then also $G_F(A)$, $G_v(A)$ and A are domains.*

Proof. By Corollary 4.3 we get that $G_v(G_F(A))$ is a domain. Since all principal symbol maps involved are multiplicative, we obtain that $G_F(A)$ is a domain, hence A is a domain too. From $G_f(G_v(A)) = G_v(G_F(A))$ we get that $G_f(G_v(A))$ is a domain and thus we deduce that $G_v(A)$ is a domain. $\square$

Now the problem of finding an extending non-commutative valuation is reduced to finding a reductor in an associated graded ring having a domain for its reduction.

In the following we show that in case the algebras are given by a finite number of relations we may relate the existence of a reductor to properties of good reduction. Let A be a filtered K -algebra with a finite filtration FA , $n = \dim_K F_1 A$ and assume that $A = K[F_1 A]$. Then we may give A as an epimorphic image of $\mathcal{F} = K \langle \underline{X} \rangle$,

the free K -algebra on the set $\underline{X} = \{X_1, \dots, X_n\}$. Say $\{x_1, \dots, x_n\}$ is a K -basis for F_1A and $\pi : K \langle \underline{X} \rangle \rightarrow A$ the canonical K -algebras morphism given by $\pi(X_i) = x_i$, $i = 1, \dots, n$. The filtration we use on $\mathcal{F}$ is the degree filtration (the total degree in the $X_1, \dots, X_n$). It follows that π is a *strict* filtered morphism, i.e. $F_n\mathcal{F}$ is taken exactly to F_nA by π . Denote by $\mathcal{R}$ the ideal of relations of A , that is $\mathcal{R} = \ker \pi$. Then

$$0 \longrightarrow \mathcal{R} \longrightarrow \mathcal{F} \longrightarrow A \longrightarrow 0 \quad (2)$$

is a strict exact sequence and therefore we obtain an exact sequence of graded $\mathcal{F}$ -modules $0 \longrightarrow \tilde{\mathcal{R}} \longrightarrow \tilde{\mathcal{F}} \longrightarrow \tilde{A} \longrightarrow 0$ by passing to the Rees objects; see Li and Van Oystaeyen (1995). Also from strict exactness of sequence (2) it follows that $G_{\mathcal{F}}(A) = G_F(R)$, where $G_{\mathcal{F}}(A)$ is the associated graded ring of A as a filtered $\mathcal{F}$ -module. Thus the sequence $0 \longrightarrow G(\mathcal{R}) \longrightarrow G(\mathcal{F}) \longrightarrow G(A) \longrightarrow 0$ is exact in $G(\mathcal{F})$ -gr. Since the filtration on $\mathcal{F}$ stems from its gradation it follows that $G(\mathcal{F}) \simeq \mathcal{F}$ and under this isomorphism $G(\mathcal{R})$ corresponds to the ideal $\mathcal{R}^\bullet$ in $\mathcal{F}$, where $\mathcal{R}^\bullet$ is the graded ideal generated by the homogeneous components of highest degrees of elements of $\mathcal{R}$.

Let us recall Theorem 2.13 from Hussein and Van Oystaeyen (1996).

Theorem 4.5. *Let A be as before. If $G_F(A)$ reduces well at O_v , then $G_F(A)$ is isomorphic to $\mathcal{F}/\mathcal{R}^\bullet$ where $\mathcal{R}$ is generated as a two-sided ideal by a finite set $\{p_1(\underline{X}), \dots, p_d(\underline{X})\}$ of elements of $\mathcal{F}$ having as homogeneous parts of highest degree $q_1(\underline{X}), \dots, q_d(\underline{X})$ which generate $\mathcal{R}^\bullet$ as a two-sided ideal. Moreover A reduces well at O_v .*

Proposition 4.1 and Theorem 4.5 together provide complete information concerning lifting properties from $G_F(A)$ to A with respect to the existence of valuation rings in $Q_{cl}(A)$ (or on some microlocalization in case non-Noetherian or non-Goldie rings are being considered) extending O_v . Note that for $\Gamma = \mathbb{Z}$ Theorem 4.5 allows ramification of valuations while the reductor property involved in Proposition 4.1 provides only unramified extensions of valuations. The latter is the case in $D(g) = Q_{cl}(g)$, Weyl algebras (see Van Oystaeyen and Willaert, 1996), Sklyanin algebras (see Hussein and Van Oystaeyen, 1996) and many more. The generalized gauge algebras provide an example for the first case.

ACKNOWLEDGMENTS

This research was supported by the bilateral project ‘‘Hopf Algebras in Algebra, Topology, Geometry and Physics’’ of the Flemish and Romanian governments. The first author gratefully acknowledges the support of the University of Antwerp.

REFERENCES

- [1] Berger, R. (1992). The quantum Poincaré-Birkhoff-Witt theorem. *Comm. Math. Phys.* 143:215–234.
- [2] Fossum, R. (1973). *The divisor class group of a Krull domain*. New York: Springer.
- [3] Van Geel, J. (1981). *Places and valuations in non-commutative ring theory*. Vol. 71. Lecture Notes in Mathematics, M. Dekker.
- [4] Hussein, M., Van Oystaeyen, F. (1996). Discrete valuations extend to certain algebras of quantum type. *Comm. Alg.* 24:2551–2566.

- [5] Li, H., Van Oystaeyen, F. (1990). Filtrations on simple artinian rings. *J. Alg.* 132:361–376.
- [6] Li, H., Van Oystaeyen, F. (1995). *Zariskian filtrations*. Vol. 2. Kluwer Monographs, Kluwer Acad. Publ.
- [7] Van Oystaeyen, F. (1975). *Prime spectra in non-commutative algebra*. Vol. 444. Lecture Notes in Mathematics, Springer.
- [8] Van Oystaeyen, F. (2000). *Algebraic geometry for associative algebras*. Vol. 232. Monographs and Textbooks in Pure and Applied Mathematics, Marcel Dekker.
- [9] Van Oystaeyen, F., Willaert, L. (1996). Valuations on extensions of Weyl skewfields. *J. Alg.* 183:359–364.
- [10] Willaert, L. (1997). Discrete valuations on Weyl skewfields. *J. Alg.* 187:537–547.